

8-2008

Towards a More Adequate Use of Interval-Valued Fuzzy Techniques in Intelligent Control: A Fuzzy Analogue of Unimodality

Van Nam Huynh

Vladik Kreinovich

The University of Texas at El Paso, vladik@utep.edu

Follow this and additional works at: https://scholarworks.utep.edu/cs_techrep



Part of the [Computer Engineering Commons](#)

Comments:

Technical Report: UTEP-CS-08-33

Published in the *Proceedings of the International Workshop on Soft Computing for Knowledge Technology*, in conjunction with The Tenth Pacific Rim International Conference on Artificial Intelligence PRICAI'08, Hanoi, Vietnam, December 15-19, 2008, pp. 80-89.

Recommended Citation

Huynh, Van Nam and Kreinovich, Vladik, "Towards a More Adequate Use of Interval-Valued Fuzzy Techniques in Intelligent Control: A Fuzzy Analogue of Unimodality" (2008). *Departmental Technical Reports (CS)*. 114.

https://scholarworks.utep.edu/cs_techrep/114

This Article is brought to you for free and open access by the Computer Science at ScholarWorks@UTEP. It has been accepted for inclusion in Departmental Technical Reports (CS) by an authorized administrator of ScholarWorks@UTEP. For more information, please contact lweber@utep.edu.

Towards a More Adequate Use of Interval-Valued Fuzzy Techniques in Intelligent Control: A Fuzzy Analogue of Unimodality

Van Nam Huynh¹ and Vladik Kreinovich²

¹ School of Knowledge Science
Japan Advanced Institute of Science
and Technology JAIST
1-1 Asahidai, Nomi
Ishikawa 923-1292 Japan
email huynh@jaist.ac.jp

² Department of Computer Science
University of Texas at El Paso
500 W. University
El Paso, TX 79968, USA
email vladik@utep.edu
Corresponding author

Abstract. It is known that interval-valued fuzzy sets provide a more adequate description of expert uncertainty than the more traditional “type-1” (number-valued) fuzzy techniques. In the current approaches for using interval-valued fuzzy techniques, it is usually assumed that all fuzzy sets $m(x) \in [\underline{m}(x), \overline{m}(x)]$ are possible. In this paper, we show that it is reasonable to restrict ourselves only to fuzzy numbers $m(x)$, i.e., “unimodal” fuzzy sets. We also describe feasible algorithms for implementing thus modified intelligent control.

Keywords: interval-valued fuzzy sets, fuzzy control, fuzzy numbers, unimodality, defuzzification, feasible algorithm


1 Introduction

Need for intelligent control. In many practical control situations, there is a small number of experts skilled in the corresponding control. Since there are only a few such skilled experts, they are unable to personally control all needed situations. It is therefore desirable to design an automated system that would implement their expertise.

Need to use fuzzy sets. Experts are often only able to describe their control by using imprecise (*fuzzy*) words from natural language such as “small” or “close to 0”. To translate such knowledge into a numerical strategy, Zadeh invented fuzzy logic. For each natural-language property P like “small” and for every possible value x of the corresponding quantity, an expert is often not 100% certain whether x satisfies the property P . We describe his or her certainty by a

degree $m(x)$ from the interval $[0, 1]$. These degrees form a *fuzzy set*. To be a more precise, a fuzzy set is usually defined as a function m which maps all possible values of the corresponding quantity into the interval $[0, 1]$; see, e.g., [2, 7]. The function m is also called a *membership function*.

“Unimodular” fuzzy sets – fuzzy numbers. Usually, fuzzy sets are “unimodular” in the sense that the corresponding membership function $m(x)$ first (non-strictly) increases (usually, from 0 to 1), and then (non-strictly) decreases (usually, from 1 to 0). Such fuzzy sets are also known as *fuzzy numbers*.




Need for a defuzzification. Based on the expert’s rules and the formulas of fuzzy logic, we translate the fuzzy sets corresponding to the natural-language terms into a fuzzy set that describes reasonable control values. Since we want a single control values, we must use a special *defuzzification* procedure.


Usually, a *centroid* defuzzification is used, in which we transform a membership function $m(x)$ into the “centroid” value

$$u = \frac{\int x \cdot m(x) dx}{\int m(x) dx}. \quad (1)$$

Need for interval-valued fuzzy sets. In practice, just like an expert cannot be 100% sure whether a given value x is small, this same expert cannot describe her degree of certainty by an exact number. At best, she can produce an interval $[\underline{m}(x), \overline{m}(x)]$ of possible values. As a result, we get *interval-valued* fuzzy sets; see, e.g., [5, 6].



The resulting interval-valued fuzzy set can be viewed as a class of all fuzzy sets m for which, for every x , the value $m(x)$ is within this interval:



How to defuzzify an interval-valued fuzzy set. As a result of defuzzifying an interval-valued fuzzy set $[\underline{m}(x), \overline{m}(x)]$, it is thus reasonable to take the interval formed by the results of defuzzifying all fuzzy sets $m(x) \in [\underline{m}(x), \overline{m}(x)]$.

Efficient algorithms have been designed for computing this interval; see, e.g., [3–6]. These algorithms and the main ideas behind these intervals are presented in the following text.

Problem. As we will see, the endpoints of these intervals are sometimes only attained for un-natural fuzzy sets – which are not fuzzy numbers.

What we do in this paper. In this paper, we propose to restrict ourselves to fuzzy numbers $m(x)$, and we design a feasible algorithm for computing the resulting (narrower) interval.

2 Algorithm for Defuzzification of Interval-Valued Fuzzy Sets: Reminder

Need to describe the algorithm and its motivations. Since our objective is to explain the problems with the existing defuzzification algorithm for interval-valued fuzzy sets, and to propose a modified algorithm that solves these problems, let us first describe this algorithm – and the motivations behind this algorithm.

How can we represent a generic membership function in a computer. We are interested in producing the algorithm for defuzzification. The input to this algorithm is a membership function. So, to describe the algorithm, we first need to describe how we can represent a generic membership function $m(x)$ in a computer.

Some membership functions are determined by their analytical (or algorithmic) expression. For example, a piece-wise linear membership function pictured above can be represented by explicit formulas for its linear parts. However, for a generic membership function, there is no analytical or algorithmic expressions.

Instead, from experts, we get the degrees $m(x_i)$ to which different values x_i are possible. In practice, we can only ask a finite number of questions to an expert, so we have only finitely many values x_i .

It is therefore reasonable to represent the “input” membership functions – describing such terms as “small” – by their values at a finite number of points.

Usually, a membership function $m(x)$ is represented by its values $m_i \stackrel{\text{def}}{=} m(x_i)$ on a uniform grid $x_i = x_0 + i \cdot h$ for some $h > 0$.

How to describe centroid defuzzification under this representation.

When we know the values $m_i = m(x_i)$ of a function $m(x)$ on a grid, a natural way to approximate an integral $\int m(x) dx$ of this function is by using the corresponding integral sum: $\int m(x) dx \approx \sum_{i=1}^n m(x_i) \cdot \Delta x_i$, where $\Delta x_i = x_{i+1} - x_i = h$. In other words, the resulting integral sum is simply proportional to the sum of the corresponding values: $\int m(x) dx \approx h \cdot \sum_{i=1}^n m_i$.

Similarly, the integral in the numerator of the centroid formula can be approximated as $\int x \cdot m(x) dx \approx h \cdot \sum_{i=1}^n x_i \cdot m_i$. When we divide this integral sum by the previous one, the factors h in the numerator and in the denominator cancel each other, so we end up with the following formula for the result u of centroid defuzzification:

$$u = \frac{\sum_{i=1}^n x_i \cdot m_i}{\sum_{i=1}^n m_i}. \quad (2)$$

Towards defuzzification for interval-valued fuzzy sets. In the interval-valued case, for every i , instead of the exact value of m_i , we only know the interval $[\underline{m}_i, \overline{m}_i]$ of possible values of m_i , where $\underline{m}_i \stackrel{\text{def}}{=} \underline{m}(x_i)$ and $\overline{m}_i \stackrel{\text{def}}{=} \overline{m}(x_i)$.

For different values $m_i \in [\underline{m}_i, \overline{m}_i]$, we get, in general, different values of u . Our objective is to find the range of possible value of u when $m_i \in [\underline{m}_i, \overline{m}_i]$.

The function (2) is continuous; thus, its range on a connected closed bounded box $[\underline{m}_1, \overline{m}_1] \times \dots \times [\underline{m}_n, \overline{m}_n]$ is an interval. We will denote the endpoints of this interval by \underline{u} and \overline{u} . Thus, to find the range, it is sufficient to find the smallest possible and the largest possible values of the expression (2) under the condition $m_i \in [\underline{m}_i, \overline{m}_i]$.

Derivation of the formula for \overline{u} . Let us start with the maximum. Let $\tilde{m}_1, \dots, \tilde{m}_n$ be the values at which the maximum is attained. It is well known, from calculus, that when maximum is attained inside the interval $\tilde{m}_i \in (\underline{m}_i, \overline{m}_i)$, then the corresponding partial derivative $\frac{\partial u}{\partial m_i}$ is equal to 0.

When the maximum is attained at $\tilde{m}_i = \underline{m}_i$, then we cannot have $\frac{\partial u}{\partial m_i} > 0$, since then, for some small $\varepsilon > 0$, the value at $m_i = \tilde{m}_i + \varepsilon$ will be even larger. Thus, we must have $\frac{\partial u}{\partial m_i} \leq 0$.

Similarly, when the maximum is attained at $\tilde{m}_i = \overline{m}_i$, then we cannot have $\frac{\partial u}{\partial m_i} < 0$, since then, for some small $\varepsilon > 0$, the value at $m_i = \tilde{m}_i - \varepsilon$ will be even larger. Thus, we must have $\frac{\partial u}{\partial m_i} \geq 0$.

The partial derivative of the expression (2) is straightforward to compute: it is equal to

$$\frac{\partial}{\partial m_i} \left(\frac{\sum_{j=1}^n x_j \cdot m_j}{\sum_{j=1}^n m_j} \right) = \frac{x_i \cdot \left(\sum_{j=1}^n m_j \right) - \sum_{j=1}^n x_j \cdot m_j}{\left(\sum_{j=1}^n m_j \right)^2} = \frac{x_i - \bar{u}}{\sum_{j=1}^n m_j}.$$

Since all the values of m_j of the membership function are non-negative, the sign of the partial derivative coincides with the sign of the difference $x_i - \bar{u}$.

Thus, we arrive at the following conclusions:

- if $\underline{m}_i < \tilde{m}_i < \overline{m}_i$, then $x_i = \bar{u}$;
- if $\tilde{m}_i = \underline{m}_i$, then $x_i \leq \bar{u}$;
- if $\tilde{m}_i = \overline{m}_i$, then $x_i \geq \bar{u}$.

So, if $x_i < \bar{u}$, we cannot have $\underline{m}_i < \tilde{m}_i < \overline{m}_i$ and we cannot have $\tilde{m}_i = \overline{m}_i$, so the only remaining possibility is $\tilde{m}_i = \underline{m}_i$.

Similarly, if $x_i > \bar{u}$, we cannot have $\underline{m}_i < \tilde{m}_i < \overline{m}_i$ and we cannot have $\tilde{m}_i = \underline{m}_i$, so the only remaining possibility is $\tilde{m}_i = \overline{m}_i$.

It should be mentioned that when $x_i = \bar{u}$, then replacing \tilde{m}_i with any other value $m_i \in [\underline{m}_i, \overline{m}_i]$ does not change the expression (2) and thus, for this particular i , we can pick any value $m_i \in [\underline{m}_i, \overline{m}_i]$.


Thus, we arrive at the following formula.

Resulting formula for \bar{u} . In the discrete case, the maximum \bar{u} is attained when we choose $m_i = \underline{m}_i$ for all i for which $x_i < \bar{u}$ and $m_i = \overline{m}_i$ for all i for which $x_i \geq \bar{u}$:

$$\bar{u} = \frac{\sum_{i: x_i < \bar{u}} x_i \cdot \underline{m}_i + \sum_{j: x_j \geq \bar{u}} x_j \cdot \overline{m}_j}{\sum_{i: x_i < \bar{u}} \underline{m}_i + \sum_{j: x_j \geq \bar{u}} \overline{m}_j}. \quad (3)$$

Similarly, in the continuous case, the maximum \bar{u} is attained when we choose $m(x) = \underline{m}(x)$ for all $x < \bar{u}$ and $m(x) = \overline{m}(x)$ for all $x \geq \bar{u}$:

$$\bar{u} = \frac{\int_{-\infty}^{\bar{u}} x \cdot \underline{m}(x) dx + \int_{\bar{u}}^{\infty} x \cdot \overline{m}(x) dx}{\int_{-\infty}^{\bar{u}} \underline{m}(x) dx + \int_{\bar{u}}^{\infty} \overline{m}(x) dx}. \quad (4)$$




Resulting formula for \underline{u} . Similarly, we can conclude that in the discrete case, the minimum \underline{u} is attained when we choose $m_i = \bar{m}_i$ for all i for which $x_i < \underline{u}$ and $m_i = \underline{m}_i$ for all i for which $x_i \geq \underline{u}$:

$$\underline{u} = \frac{\sum_{i:x_i < \underline{u}} x_i \cdot \bar{m}_i + \sum_{j:x_j \geq \underline{u}} x_j \cdot \underline{m}_j}{\sum_{i:x_i < \underline{u}} \bar{m}_i + \sum_{j:x_j \geq \underline{u}} \underline{m}_j}. \quad (5)$$

In the continuous case, the minimum \underline{u} is attained when we choose $m(x) = \bar{m}(x)$ for all $x < \underline{u}$ and $m(x) = \underline{m}(x)$ for all $x \geq \underline{u}$:

$$\underline{u} = \frac{\int_{-\infty}^{\underline{u}} x \cdot \bar{m}(x) dx + \int_{\underline{u}}^{\infty} x \cdot \underline{m}(x) dx}{\int_{-\infty}^{\underline{u}} \bar{m}(x) dx + \int_{\underline{u}}^{\infty} \underline{m}(x) dx}. \quad (6)$$



How can we actually compute \bar{u} and \underline{u} : analytical case. For the case when $\underline{m}(x)$ and $\bar{m}(x)$ are given by analytical formulas, we can explicitly integrate both numerator and denominator and get algebraic equations for the unknown values \bar{u} or \underline{u} .

How can we actually compute \bar{u} and \underline{u} : towards an algorithm for the general case. How can we perform these computations in the general case? The above formulas (3) and (5) require that we know \bar{u} and \underline{u} in order to find the appropriate values $m_i \in [\underline{m}_i, \bar{m}_i]$. Thus, the above formulas do not directly lead to an efficient algorithm for computing \bar{u} and \underline{u} .

The possibility to efficiently compute \bar{u} and \underline{u} comes from the fact that, e.g., in the formula (3), all we need to know is where exactly \bar{u} is in comparison with the values $x_1 < x_2 < \dots < x_n$. For simplicity, let us supplement these values with $x_0 = -\infty$ and $x_{n+1} = +\infty$. Then, the real line is divided into $n + 1$ (finite or infinite) intervals $(x_k, x_{k+1}]$, $k = 0, 1, \dots, n$. So, to find \bar{u} , it is sufficient to try all these $n + 1$ intervals.

We will describe the arguments in details for the case of the maximum. For the minimum, the arguments are similar.

If $x_k < \bar{u} \leq x_{k+1}$, then the formula (3) can be rewritten as $\bar{u} = \bar{u}_k \stackrel{\text{def}}{=} \frac{\bar{N}_k}{\bar{D}_k}$, where

$$\bar{N}_k \stackrel{\text{def}}{=} \sum_{i=1}^k x_i \cdot \underline{m}_i + \sum_{j=k+1}^n x_j \cdot \bar{m}_j,$$

and

$$\bar{D}_k \stackrel{\text{def}}{=} \sum_{i=1}^k \underline{m}_i + \sum_{j=k+1}^n \bar{m}_j.$$

We only need to consider values k for which $x_k < \bar{u}_k \leq x_{k+1}$.

So, we compute the ratios \bar{u}_k for all k , keep only those ratios for which the inequality $x_k < \bar{u}_k \leq x_{k+1}$ is satisfied, and then return the largest of the kept ratios \bar{u}_k as the desired value of \bar{u} .

Computational complexity of the resulting algorithm: discussion. How many computational steps do we need to perform these computations? For the standard defuzzification (2), we need to perform a liner number of steps $O(n)$: n multiplications and $n - 1$ additions to compute the numerator, $n - 1$ additions to compute the denominator, and 1 division to compute the ratio u . Let us show that we can compute \bar{u} in linear time as well.

For $k = 0$, we can compute \bar{N}_0 and \bar{D}_0 in linear time. Then, when we move from \bar{N}_k to \bar{N}_{k+1} (or from \bar{D}_k to \bar{D}_{k+1}), we only to change one term, so we only need a finite number of steps. Thus, to find all n ratios, we only need a linear number of steps.

Let us summarize the resulting algorithm.

Algorithm for computing \bar{u} .

- First, we compute $\bar{N}_0 = \sum_{j=1}^n x_j \cdot \bar{m}_j$ and $\bar{D}_0 = \sum_{j=1}^n \bar{m}_j$.
- Then, for $k = 1, 2, \dots, n$, we compute $\bar{N}_{k+1} = \bar{N}_k - x_k \cdot (\bar{m}_k - \underline{m}_k)$ and $\bar{D}_{k+1} = \bar{D}_k - (\bar{m}_k - \underline{m}_k)$.

- For each k , we compute the ratio $\bar{u}_k = \frac{\bar{N}_k}{\bar{D}_k}$, and check whether

$$x_k < \bar{u}_k \leq u_{k+1};$$

if this inequality is satisfied, we keep \bar{u}_k as a possible value.

- The largest of these possible values is then returned as \bar{u} .

Comment. A similar efficient (linear time) algorithm can be used to compute \underline{u} .

Algorithm for computing \underline{u} .

- First, we compute $\underline{N}_0 = \sum_{j=1}^n x_j \cdot \underline{m}_j$ and $\underline{D}_0 = \sum_{j=1}^n \underline{m}_j$.
- Then, for $k = 1, 2, \dots, n$, we compute $\underline{N}_{k+1} = \underline{N}_k + x_k \cdot (\bar{m}_k - \underline{m}_k)$ and $\underline{D}_{k+1} = \underline{D}_k + (\bar{m}_k - \underline{m}_k)$.
- For each k , we compute the ratio $\underline{u}_k = \frac{\underline{N}_k}{\underline{D}_k}$, and check whether

$$x_k < \underline{u}_k \leq u_{k+1};$$

if this inequality is satisfied, we keep \underline{u}_k as a possible value.

- The smallest of these possible values is then returned as \underline{u} .

3 Towards a More Adequate Defuzzification

The problem. The problem is that,

- as we have mentioned earlier, it is often reasonable to restrict ourselves to fuzzy numbers (unimodal fuzzy sets),
- while, as we have seen, the maximum and/or minimum of the value u is sometimes attained at a membership function which is not unimodal.

It is therefore desirable to find the maximum and the minimum of u only among *unimodal* values m_i , i.e., values which m_i first (non-strictly) increase) and then (non-strictly) decreases.

The problem reformulated in precise mathematical terms. In precise terms, we are only interested in finding the maximum and the minimum of the expression (2) among all the values m_1, \dots, m_n for which, for some “mode location” $\ell = 1, 2, \dots, n$, we have

$$m_1 \leq m_2 \leq \dots \leq m_{\ell-1} < m_{\ell} \geq m_{\ell+1} \geq \dots \geq m_{n-1} \geq m_n.$$

Let us denote the corresponding minimum and maximum by \underline{u}_a and \bar{u}_a (where a stands for “adequate”).

Towards a solution of the problem. Let us fix the values \bar{u}_a and ℓ and see how we can use the inequalities corresponding to this value.

When $x_i < \bar{u}_a$, then, as we mentioned earlier, the derivative $\frac{\partial u}{\partial m_i}$ is negative and thus, we cannot decrease \tilde{m}_i . In the past, we only had one restriction: that $m_i \geq \underline{m}_i$. Now, we have additional restrictions: e.g., for $i \leq \ell$, that $m_i \geq m_j$ for all $j < i$. Thus, the fact that we cannot decrease m_i means that either $\tilde{m}_i = \underline{m}_i$ or that $\tilde{m}_i = \tilde{m}_j$ for some $j < i$. In the second case, for \tilde{m}_j , we can repeat the same argument, and eventually, we will find that $\tilde{m}_i = \tilde{m}_j$ for some value j which cannot be decreased because it is equal to $\tilde{m}_j = \underline{m}_j$. Thus, we have $\tilde{m}_i = \underline{m}_j$.

In general, since $i \leq \ell$, we have $\tilde{m}_i \geq \tilde{m}_j \geq \underline{m}_j$. Thus, we have $\tilde{m}_i \geq \max(\underline{m}_1, \dots, \underline{m}_i)$. Since we concluded that \tilde{m}_i is equal to one of these lower endpoints, it cannot be larger than the largest of them, so we have $\tilde{m}_i = \max(\underline{m}_1, \dots, \underline{m}_i)$.

For $i > \ell$, we may also have $\tilde{m}_i = \bar{m}_j$ for some j for which $x_j > \bar{v}_a$. In this case, the values m_k between i and j are constant.

Thus, the “past-mode” part ($i > \ell$) of the optimal solution can be divided into three zones:

- first, there is a zone $[\ell, s]$ (s for *start*) before \bar{v}_a where we have


$$\tilde{m}_i = \max(\underline{m}_i, \dots, \underline{m}_s);$$

- then, there is a zone $[e, n]$ (e for *end*) past \bar{v}_a where we have

$$\tilde{m}_i = \min(\bar{m}_i, \dots, \bar{m}_n);$$

- finally, in the zone between s and e , the values are constant.

So, to describe all such solutions, it is sufficient to try all possible values of three indices: ℓ , s and e .



Resulting algorithm for computing \bar{u}_a .

- First, for all i and j , we compute $m_{ij}^- \stackrel{\text{def}}{=} \max(\underline{m}_i, \dots, \underline{m}_j)$ and $m_{ij}^+ \stackrel{\text{def}}{=} \min(\bar{m}_i, \dots, \bar{m}_j)$. For each i , computing the next value m_{ij}^\pm from the previous one requires one step, all these values can be computed in time $O(n^2)$.

- Second, for each of n^3 possible combinations of three integers $\ell \leq s < e$, we take $m_i = m_{1i}^-$ for $i < \ell$, $m_i = m_{is}^-$ for $\ell \leq i \leq s$, $m_i = m_{in}^+$ for $i \geq e$, and $m_i = \text{const} \in [m_e, m_s]$ for $i \in (e, s)$.
- We check whether all these values satisfy the conditions $m_i \in [\underline{m}_i, \overline{m}_i]$, and if yes, we compute the ratio u .
- The largest of these values is returned as the desired upper bound \bar{u}_a .

Computational complexity. For each of $O(n^3)$ combinations of values, we need linear time to compute the ratio u . Thus, totally, we need $O(n^3) \cdot O(n) = O(n^4)$ steps. This is still polynomial time, i.e., this algorithm is still feasible; see, e.g., [1].

Comment. A similar algorithm can be described for computing \underline{u}_a .

Algorithm for computing \underline{u}_a .

- For each of n^3 possible combinations of three integers $s \leq e \leq \ell$, we take $m_i = m_{in}^-$ for $i > \ell$, $m_i = m_{si}^-$ for $s \leq i \leq \ell$, $m_i = m_{1i}^+$ for $i \leq s$, and $m_i \in [m_e, m_s]$ for $i \in (e, s)$.
- We check whether all these values satisfy the conditions $m_i \in [\underline{m}_i, \overline{m}_i]$, and if yes, we compute the ratio u .
- The smallest of these values is returned as the desired lower bound \underline{u}_a .

Acknowledgments

This work was supported in part by NSF grant HRD-0734825, by Grant 1 T36 GM078000-01 from the National Institutes of Health, and by the Japan Advanced Institute of Science and Technology (JAIST) International Joint Research Grant 2006-08.

References

1. Cormen, T. H., Leiserson, C. E., Rivest, R. L., Stein, C.: *Introduction to Algorithms*, MIT Press, Cambridge, Massachusetts, 2001.
2. Klir, G., Yuan, B.: *Fuzzy Sets and Fuzzy Logic*, Prentice Hall, New Jersey, 1995.
3. Kreinovich, V., Lakeyev, A., Rohn, J., Kahl, P.: *Computational Complexity and Feasibility of Data Processing and Interval Computations*, Kluwer Academic Press, 1998.
4. Lea, R., Kreinovich, V., Trejo, R.: “Optimal interval enclosures for fractionally-linear functions, and their application to intelligent control”, *Reliable Computing* 2(3):265–286, 1996.
5. Mendel, J. M.: *Uncertain Rule-Based Fuzzy Logic Systems: Introduction and New Directions*, Prentice-Hall, 2001.
6. Mendel, J. M.: “Type-2 fuzzy sets and systems: an overview”, *IEEE Computational Intelligence Magazine* 2:20–29, 2007.
7. Nguyen, H. T., Walker, E. A.: *First Course in Fuzzy Logic*, CRC Press, Boca Raton, Florida, 2006.