

From $[0,1]$ -Based Logic To Interval Logic

(From known description of all possible
 $[0,1]$ -based logical operations
to a description of all possible
interval-based logical operations)

Hung T. Nguyen¹ and Vladik Kreinovich²

¹Department of Mathematical Sciences
New Mexico State University
Las Cruces, NM 88003, USA
hunguyen@nmsu.edu

²Department of Computer Science
University of Texas at El Paso
El Paso, TX 79968, USA
vladik@cs.utep.edu

Abstract

Since early 1960s, we have a complete description of all possible $[0, 1]$ -based logical operations, namely of “and”-operations (t-norms) and of “or”-operations (t-conorms). In some real-life situations, intervals provide a more adequate way of describing uncertainty, so we need to describe interval-based logical operations. Usually, researchers followed a pragmatic path and simply derived these operations from the $[0, 1]$ -based ones. From the foundational viewpoint, it is desirable not to *a priori* restrict ourselves to such derivative operations but, instead, to get a description of all interval-based operations which satisfy reasonable properties.

Such description is presented in this paper. It turns out that all such operations can be described as the result of applying interval computations to the corresponding $[0, 1]$ -based ones.

1 [0,1]-Based Logical Operations: Reminder

1.1 Why [0,1]-Based Logical Values

In many areas of expertise, such as medicine, geophysics, etc., human experts are needed. Usually, there are very few top level experts, and it is not physically possible for these few experts to solve all numerous related problems. It is therefore desirable to develop a computer-based system which incorporates the knowledge of the top experts and uses this knowledge either to directly solve the related problems – or, at least, to provide high-level advice to people trying to solve these problems.

Experts can describe their knowledge in terms of statements and rules, but this formulation often comes with uncertainty and ambiguity: experts are often not 100% confident in the statements which form their knowledge, and even when they are, these statements are formulated in terms of words of natural language (such as “large”) which do not have precise meaning. To adequately describe the expert knowledge, we must therefore store, in the knowledge base, not only the statements themselves, but also the indication of the degree to which the experts are confident in these statements.

This degree is usually described by a number from the interval $[0,1]$. An expert’s degree of confidence $d(A)$ in a statement A can be determined, if, e.g., we ask an expert to estimate his/her degree of confidence on a scale from 0 to 10. If s/he selects 8, then we take $d(A) = 8/10$.

1.2 Why [0,1]-Based Logical Operations

Suppose now that we know the degrees of confidence $d(A)$ and $d(B)$ in statements A and B , and we know nothing else about A and B . Suppose also that we are interested in the degree of confidence of the composite statement $A \& B$. Since the only information available consists of the values $d(A)$ and $d(B)$, we must compute $d(A \& B)$ based on these values. We must be able to do that for arbitrary values $d(A)$ and $d(B)$. Therefore, we need a *function* that transforms the values $d(A)$ and $d(B)$ into an estimate for $d(A \& B)$. Such a function is called an “*and*”-operation (*t-norm*). If an “and”-operation $f_{\&} : [0, 1] \times [0, 1] \rightarrow [0, 1]$ is fixed, then we take $f_{\&}(d(A), d(B))$ as an estimate for $d(A \& B)$.

Similarly, to estimate the degree of confidence in $A \vee B$, we need an “*or*”-operation (*t-conorm*) $f_{\vee} : [0, 1] \times [0, 1] \rightarrow [0, 1]$. The following are the natural *general* requirements for “and”- and “or”-operations:

Definition 1.

- By an “and”-operation, we mean a commutative, associative, monotonic, continuous operation $f_{\&} : [0, 1] \times [0, 1] \rightarrow [0, 1]$ for which $f_{\&}(1, a) = a$ and $f_{\&}(0, a) = 0$.
- By an “or”-operation, we mean a commutative, associative, monotonic, continuous operation $f_{\vee} : [0, 1] \times [0, 1] \rightarrow [0, 1]$ for which $f_{\vee}(1, a) = 1$ and $f_{\vee}(0, a) = a$.

These properties are easy to explain. For example, commutativity $f_{\&}(a, b) = f_{\&}(b, a)$ comes from the fact that, from a common sense viewpoint, composite statements $A \& B$ and $B \& A$ are equivalent; therefore, we expect our “and”-operation to lead to the same degree of certainty for both composite statements. In precise terms, this means that we expect $f_{\&}(d(A), d(B)) = f_{\&}(d(B), d(A))$ for every two statements A and B . If we denote $d(A)$ by a and $d(B)$ by b , we can therefore conclude that $f_{\&}(a, b) = f_{\&}(b, a)$ for every a and b .

Similarly, associativity $f_{\&}(a, f_{\&}(b, c)) = f_{\&}(f_{\&}(a, b), c)$ comes from the fact that from the common sense viewpoint, the composite statements $A \& (B \& C)$ and $(A \& B) \& C$ are equivalent.

Monotonicity, i.e., the fact that $a_1 \leq a_2$ and $b_1 \leq b_2$, then $f_{\&}(a_1, b_1) \leq f_{\&}(a_2, b_2)$ (and $f_{\vee}(a_1, b_1) \leq f_{\vee}(a_2, b_2)$), comes from the fact that if our degree of confidence in A_1 is smaller than the degree of confidence in A_2 , and the degree of confidence in B_1 is smaller than the degree of confidence in B_2 , then our confidence in $A_1 \& B_1$ must be smaller (or at least equal, but not larger) than our confidence in $A_2 \& B_2$.

The first two pairs of “and” and “or” operations were proposed by L. Zadeh in [28]: $f_{\&}(x, y) = \min(x, y)$, $f_{\vee}(x, y) = \max(x, y)$, and $f_{\&}(x, y) = x \cdot y$, $f_{\vee}(x, y) = x + y - x \cdot y$. Later, numerous other operations have been proposed: e.g., in [8], Giles proposed “bold and” $f_{\&}(a, b) = \max(a + b - 1, 0)$ and “bold or” $f_{\vee}(a, b) = \min(a + b, 1)$.

1.3 Fuzzy Control: One of the Main Applications of Fuzzy Logic

One of the main applications of fuzzy logic is *fuzzy control* (see, e.g., [17]). In most industrial applications, we want to control the corresponding industrial processes in such a way as to maximize the output within certain (physical and economical) restrictions. When the corresponding mathematical description is linear, we can use well-known optimal control techniques to find the optimal control strategy. In reality, however, most industrial processes are non-linear. For non-linear control problems, the situation is much more complicated: there are good recipes which often work but, alas, there is still no general method of generating an optimal (or even a reasonably good) control.

If for a certain industrial process, no known technique leads to a good quality control, what can we do? Usually, the very fact that this process is actually used in industry means that this process is reasonably well controlled by human controllers. Therefore, if we want to automate this control, we must somehow transform the knowledge of these expert controllers (operators) into an automatic control strategy.

Specifically, our goal is to describe a function which takes the sensor inputs x_1, \dots, x_n (numbers) and generates the (numerical) value of the control effort u . Unfortunately, expert operators cannot formulate their expertise in these terms. Instead, they describe their control strategy by using uncertain (“fuzzy”) statements of the type “if the obstacle is straight ahead, the distance to it is small, and the velocity of the car is medium, press the brakes hard”. Fuzzy control is a methodology which translates such statements into precise formulas for control.

Once we have selected a fuzzy “and”-operation $f_{\&}(a, b)$ and a fuzzy “or”-operation $f_{\vee}(a, b)$, we are able to transform an arbitrary set of fuzzy if-then rules connecting inputs x_1, \dots, x_n and the output u into a crisp function $y = f(x_1, \dots, x_n)$. Indeed, let us assume that the relation between the inputs x_1, \dots, x_n and the output u can be characterized by several if-then rules:

$$\begin{aligned}
 & \text{if } A_{11}(x_1) \text{ and } \dots \text{ and } A_{1n}(x_n) \text{ then } B_1(u); \\
 & \qquad \qquad \qquad \dots \\
 & \text{if } A_{i1}(x_1) \text{ and } \dots \text{ and } A_{in}(x_n) \text{ then } B_i(u); \qquad (1) \\
 & \qquad \qquad \qquad \dots \\
 & \text{if } A_{m1}(x_1) \text{ and } \dots \text{ and } A_{mn}(x_n) \text{ then } B_m(u),
 \end{aligned}$$

where $A_{ij}(x_j)$ and $B_i(u)$ are properties expressed by words from natural language. This interpretation consists of the following steps (see, e.g., [12, 17]):

- First, we can use one of the known elicitation techniques to determine the membership functions $\mu_{ij}^A(x_j)$ and $\mu_i^B(y)$ corresponding to the words $A_{ij}(x_j)$ and $B_i(y)$.
- Then, we can use the fuzzy “and” operation $f_{\&}(a, b)$ to determine, for each rule i , for given input x_1, \dots, x_n , and for a given control u , the degree $c_i(u)$ to which the given input and control satisfies this rule. This value is equal to

$$c_i(u) = f_{\&}(\mu_{i1}^A(x_1), \dots, \mu_{in}^A(x_n), \mu_i^B(u)). \quad (2)$$

- Next, we use the fuzzy “or” operation $f_{\vee}(a, b)$ to determine the degree $\mu(u)$ to which one of these rules is activated:

$$\mu(u) = f_{\vee}(c_1(u), \dots, c_m(u)). \quad (3)$$

- Finally, we apply one of the many known defuzzification procedures – e.g., the centroid defuzzification

$$\bar{u} = \frac{\int u \cdot \mu(u) \, du}{\int \mu(u) \, du} \quad (4)$$

– to determine the actual control value \bar{u} which we want to apply for the given input x_1, \dots, x_n .

1.4 From This Viewpoint, the More Logical Operations We Can Find, the Better

For each pair of the “and”- and “or”-operations, we can have a reasonable fuzzy control strategy. However, the fact that we have three pairs of operations does not necessarily mean that we should not look for more. It is a general commonsense fact that the more options one has, the better option one can find for some future optimization problem. This general fact is also true for fuzzy control (and for other applications of fuzzy logic). Indeed, as we have mentioned, Zadeh [28] originally proposed two pairs of operations:

- $f_{\&}(a, b) = \min(a, b)$, $f_{\vee}(a, b) = \max(a, b)$;
- $f_{\&}(a, b) = a \cdot b$, $f_{\vee}(a, b) = a + b - a \cdot b$.

For some optimality criteria, these pairs are indeed the best; for example:

- the first pair is the best when we are interested in the operations which are the most robust (the least sensitive) in the worst case;
- the second pair is the best when we are interested in the operations which are the most robust (the least sensitive) in the average.

However, for other criteria, other pairs are optimal; for example:

- when we want to achieve the most stable fuzzy control, we should use $f_{\&}(a, b) = \min(a, b)$ and $f_{\vee}(a, b) = \min(a + b, 1)$;
- when we want to achieve the most smooth fuzzy control, we should use $f_{\&}(a, b) = \max(a + b - 1, 0)$ and $f_{\vee}(a, b) = \max(a, b)$;

(for exact formulations and similar results, see, e.g., [22], the surveys [13, 17] and references therein).

1.5 Description of All Possible [0,1]-Based Logical Operations: Reminder

As we have just argued, it is desirable not to *a priori* restrict ourselves to known operations, but, instead, to get a complete description of all possible operations. For [0, 1]-based logical operations, such a classification is known.

This classification is related to the known fact that we can get new t-norms if we consider different “scales” on the interval $[0, 1]$ of all possible degrees of certainty. Namely, the assignment of different numerical degrees to words expressing uncertainty is rather arbitrary. Let us assume that we assign new values to these words, and let $\varphi(a)$ be a new value assigned to the word to which we originally assigned the value a . In this new scale, to each statement A , instead of the original degree of certainty $d(A)$, we assign a new degree of certainty $d'(A) = \varphi(d(A))$. In the new scale, the same “and”-operation will look different. Namely, if we know the degrees $a' = d'(A)$ and $b' = d'(B)$ in the new scale, and we want to find $d'(A \& B)$, then we must do the following:

- first, we compute the degrees $a = d(A)$ and $b = d(B)$ in the old scale as $a = \varphi^{-1}(a')$ and $b = \varphi^{-1}(b')$ (where φ^{-1} denotes the inverse function);
- second, we use the known t-norm $f_{\&}(a, b)$ to compute the degree of certainty $c = f_{\&}(a, b) = f_{\&}(\varphi^{-1}(a'), \varphi^{-1}(b'))$ of the composite statement $A \& B$ in the old scale;
- finally, we transform the degree c back into the new scale, resulting in $c' = \varphi(c) = \varphi(f_{\&}(\varphi^{-1}(a'), \varphi^{-1}(b')))$.

This three-step procedure is equivalent to using an operation $f'_{\&}(a', b') = \varphi(f_{\&}(\varphi^{-1}(a'), \varphi^{-1}(b')))$ and the new operation is called *isomorphic* to the original t-norm $f_{\&}(a, b)$. Isomorphic operations provide numerous new *examples* of t-norms and t-conorms.

The complete description of all possible $[0, 1]$ -based logical operations (which uses rescaling and isomorphisms) has been given, in effect, in [16] (see also [12, 14, 19, 21]). It turns out that every t-norm $f_{\&}(a, b)$ can be described as follows:

- we subdivide the interval $[0, 1]$ into subintervals;
- the restriction of the t-norm $f_{\&}(a, b)$ to each of these subintervals is isomorphic either to the “algebraic” t-norm $a \cdot b$, or to $\max(a + b - 1, 0)$, or to $\min(a, b)$; this describes the values of $f_{\&}(a, b)$ for the case when both a and b belong to the same subinterval;
- when a and b belong to different subintervals, then $f_{\&}(a, b) = \min(a, b)$.

A similar description is known for t-conorms.

2 Interval-Based Logical Operations: Reminder and Formulation of a Problem

2.1 Need for Interval-Based Logical Values

Experts cannot describe their degrees of confidence precisely. At best, they can give an *interval* of possible values. For example, an expert can point to 8 on a scale from 0 to 10, but this same expert will hardly be able to pinpoint a value on a scale from 0 to 100. As a result, the only thing that we know about the expert's degree of confidence is that it is closer to 8 than to 7 or to 9, i.e., that it is in the interval $[0.75, 0.85]$.

So, to describe degrees of confidence more adequately, we must use *intervals* $\mathbf{a} = [a^-, a^+]$ instead of real numbers. In this representation, real numbers can be viewed as particular – degenerate – cases of intervals $[a, a]$. The idea of using intervals was originally proposed by Zadeh himself and further developed by Bandler and Kohout [1], Türkşen [24], and others; for a recent survey, see, e.g., [18] (see also [2, 3, 4, 5, 6]).

2.2 Need for Interval-Based Logical Operations

Since we went from numbers to intervals in our description of degrees of certainty, we must have “and” and “or” operations as functions from intervals to intervals. For example, in fuzzy control, if the expert controller's degrees of certainty in the properties $A_{ij}(x_j)$ and $B_i(u)$ (like “small”) are described by intervals, we need operations on intervals to combine these degrees and to generate the resulting control value.

2.3 Currently Used Interval-Based Logical Operations: Reminder

Traditionally, researchers followed a pragmatic path and simply derived these operations from the $[0, 1]$ -based ones. Namely, when an expert says that his/her degree of certainty in a statement A belongs to the interval $[a^-, a^+]$, we can interpret it as meaning that the (unknown) actual degree of confidence can be any number from this interval. With this interpretation in mind, it is natural to define, e.g., an interval “and”-operation as follows:

- First, we select a $[0, 1]$ -based “and”-operation (t-norm) $f_{\&}(a, b)$. This operation corresponds to the case when an expert knows the exact values of his/her degrees of certainty, i.e., when the intervals $\mathbf{a} = [a^-, a^+]$ and $\mathbf{b} = [b^-, b^+]$ are degenerate ($a^- = a^+$ and $b^- = b^+$).
- Next, when we know the interval degrees \mathbf{a} and \mathbf{b} , we interpret these intervals by saying that a can take any value from \mathbf{a} and b can take any

value from \mathbf{b} . Thus, as the degree corresponding to $A \& B$, it is natural to take the set of all possible values of $f_{\&}(a, b)$ when $a \in \mathbf{a}$ and $b \in \mathbf{b}$. In precise terms, we define $f_{\&}(\mathbf{a}, \mathbf{b})$ as follows:

$$f_{\&}(\mathbf{a}, \mathbf{b}) = \{f_{\&}(a, b) \mid a \in \mathbf{a}, b \in \mathbf{b}\}.$$

This formula is a particular case of the so-called *interval computations* [9, 10, 11, 20]. Since the function $f_{\&}(a, b)$ is monotonically increasing and continuous, the resulting set is easy to describe:

$$f_{\&}([a^-, a^+], [b^-, b^+]) = [f_{\&}(a^-, b^-), f_{\&}(a^+, b^+)].$$

We can use a similar “pragmatic” approach and define an interval-based “or” operation as

$$f_{\vee}([a^-, a^+], [b^-, b^+]) = [f_{\vee}(a^-, b^-), f_{\vee}(a^+, b^+)].$$

For example, if we start with $f_{\&}(a, b) = \min(a, b)$ and $f_{\vee}(a, b) = \max(a, b)$, we get interval operations

$$f_{\&}([a^-, a^+], [b^-, b^+]) = [\min(a^-, b^-), \min(a^+, b^+)];$$

$$f_{\vee}([a^-, a^+], [b^-, b^+]) = [\max(a^-, b^-), \max(a^+, b^+)].$$

When we start with $f_{\&}(a, b) = a \cdot b$ and $f_{\vee}(a, b) = a + b - a \cdot b$, we get the following interval operations:

$$f_{\&}([a^-, a^+], [b^-, b^+]) = [a^- \cdot b^-, a^+ \cdot b^+];$$

$$f_{\vee}([a^-, a^+], [b^-, b^+]) = [a^- + b^- - a^- \cdot b^-, a^+ + b^+ - a^+ \cdot b^+].$$

In the general case, the resulting interval operations satisfy the same natural properties of associativity, commutativity, etc. as the original $[0, 1]$ -based ones (see, e.g., [19]).

2.4 Need for a Description of All Possible Interval-Based Logical Operations: Reminder

As we have already mentioned when we described $[0, 1]$ -based operations, the fact that we have a class of operations does not necessarily mean that we should not look for more – because the more options one has, the better option one can find for some future optimization problem. From this commonsense viewpoint, it is desirable not to *a priori* restrict ourselves to such “derivative” interval operative but, instead, to get a complete description of all possible interval-based operations.

2.5 Description of All Possible Interval-Based Logical Operations: What Was Known and What We Prove in This Paper

The task of obtaining a description of all possible interval-based logical operations was started in a pioneer paper by Zuo [29] who described all interval-based operations which are *strictly monotonic* (in some reasonable sense). In this paper, we extend Zuo's results and find a description of *all possible* interval-based logical operations (which satisfy reasonable properties like commutativity and monotonicity).

Specifically, we show that the above interval-computation operations are the only ones possible. Thus, we provide a fundamental justification for the traditional (interval-computation) approach.

3 Towards Formalization of the Problem: How to Define Monotonicity for Interval Operations?

3.1 Why Is It Important to Define Monotonicity?

An important part of the definition of t-norm and t-conorm is the requirement that these operations are *monotonic*, i.e., that if $a_1 \leq a_2$ and $b_1 \leq b_2$, then $f_{\&}(a_1, b_1) \leq f_{\&}(a_2, b_2)$ and $f_{\vee}(a_1, b_1) \leq f_{\vee}(a_2, b_2)$. For $[0, 1]$ -based operations, these properties are easy to formalize, because the order \leq is well defined on the interval $[0, 1]$. For interval degrees, however, the situation is less clear.

If we know the interval degrees $\mathbf{a}_1 = [a_1^-, a_1^+]$ and $\mathbf{a}_2 = [a_2^-, a_2^+]$ for two statements A_1 and A_2 , this means that the actual degree of confidence a_1 in A_1 can take any value from the interval \mathbf{a}_1 , and the actual degree of confidence a_2 in A_2 can take any value from the interval \mathbf{a}_2 . If the intervals \mathbf{a}_1 and \mathbf{a}_2 intersect, then, depending on the selection of the values $a_i \in \mathbf{a}_i$, we may have $a_1 < a_2$ and we may also have $a_2 < a_1$.

For example, if $\mathbf{a}_1 = [0.7, 0.9]$ and $\mathbf{a}_2 = [0.8, 1.0]$, then:

- on one hand, we may have $a_1 = 0.7 \in \mathbf{a}_1$ and $a_2 = 1.0 \in \mathbf{a}_2$, in which case $a_1 < a_2$;
- on the other hand, we may have $a_1 = 0.9 \in \mathbf{a}_1$ and $a_2 = 0.8 \in \mathbf{a}_2$, in which case $a_1 > a_2$.

Comment. A reader may notice the similarity between this example and problems from constraint propagation (see, e.g., [15, 23, 25, 26]). To get a better understanding of our problem, let us explicitly describe the similarity and the difference between similar problems from constraint propagation and the problems considered in this section.

A typical related constraint propagation problem would be formulated as follows. Suppose that in addition to the domains $\mathbf{a}_1 = [0.7, 0.9]$ and $\mathbf{a}_2 = [0.8, 1.0]$ of two quantities a_1 and a_2 , we know that $a_1 \geq a_2$. We can say that we have three *constraints*:

- the domain $\mathbf{a}_1 = [0.7, 0.9]$ is a constraint on the value of the first quantity a_1 ; it means that the value a_1 must satisfy the inequality $0.7 \leq a_1 \leq 0.9$;
- the domain $\mathbf{a}_2 = [0.8, 1.0]$ is a constraint on the value of the second quantity a_2 ; it means that the value a_2 must satisfy the inequality $0.8 \leq a_2 \leq 1.0$;
- finally, the inequality $a_1 \geq a_2$ is a joint constraint which relates the values of both variables.

These constraints lead to the updating of the previously known constraints and to the appearance of the new constraints. The corresponding process of updating previously known constraints and of discovering new (derivative) constraints is called *constraint propagation*. In the above example, we can use the three constraints to update a constraint on a_2 , namely, to deduce a stricter bound on a_2 : Indeed, since $a_2 \leq a_1$, and $a_1 \leq 0.9$, we can conclude that $a_2 \leq 0.9$ and hence, a_2 can only take values from a (narrower) interval $[0.8, 0.9]$. In more general terms:

- in constraint propagation:
 - we have intervals of possible values of certain quantities, and
 - we *know the relations* between the values of these quantities
 (and we use these relations to narrow down the intervals).
- In this section, we consider the “inverse” problem:
 - we know the intervals of possible values of certain quantities, and
 - we *want to find the relations* between the values of these quantities.

3.2 Solution: Operations “Necessarily \leq ” and “Possibly \leq ”

We have already mentioned that for interval degrees \mathbf{a}_1 and \mathbf{a}_2 , it is sometimes not clear whether $\mathbf{a}_1 \leq \mathbf{a}_2$ or not. However, the situation is not hopeless: we have the following two natural order-like relations:

Definition 2. Let $\mathbf{a}_1 = [a_1^-, a_1^+]$ and $\mathbf{a}_2 = [a_2^-, a_2^+]$ be two intervals.

- We say that \mathbf{a}_1 is necessarily \leq \mathbf{a}_2 (and denote it by $\mathbf{a}_1 \leq^\square \mathbf{a}_2$) if $a_1 \leq a_2$ for every $a_1 \in \mathbf{a}_1$ and for every $a_2 \in \mathbf{a}_2$.
- We say that \mathbf{a}_1 is possibly \leq \mathbf{a}_2 (and denote it by $\mathbf{a}_1 \leq^\diamond \mathbf{a}_2$) if $a_1 \leq a_2$ for some $a_1 \in \mathbf{a}_1$ and $a_2 \in \mathbf{a}_2$.

It is therefore natural to require that the desired interval-based logical operations be monotonic relative to *both* these operations, i.e., that:

- if $\mathbf{a}_1 \leq^\square \mathbf{a}_2$ and $\mathbf{b}_1 \leq^\square \mathbf{b}_2$, then $f_{\&}(\mathbf{a}_1, \mathbf{b}_1) \leq^\square f_{\&}(\mathbf{a}_2, \mathbf{b}_2)$;
- if $\mathbf{a}_1 \leq^\diamond \mathbf{a}_2$ and $\mathbf{b}_1 \leq^\diamond \mathbf{b}_2$, then $f_{\&}(\mathbf{a}_1, \mathbf{b}_1) \leq^\diamond f_{\&}(\mathbf{a}_2, \mathbf{b}_2)$.

We can describe these new monotonicity requirements in general terms:

Definition 3. Let L be an arbitrary (partially) ordered set.

- Let a^- and a^+ be two points from L for which $a^- \leq a^+$. The set

$$\{b \mid a^- \leq b \leq a^+\}$$

will be called an interval and denoted by $[a^-, a^+]$.

- The set of all intervals over L will be denoted by $\mathbb{I}(L)$.

For intervals over an arbitrary ordered set L , we can use Definition 2 to define relations \leq^\square and \leq^\diamond . Proposition 1 holds for this case as well. Monotonicity with respect to these operations can then be defined as follows:

Definition 4. Let n be an arbitrary positive integer.

- We say that an n -ary interval operation $F : \mathbb{I}(L) \times \dots \times \mathbb{I}(L) \rightarrow \mathbb{I}(L)$ is \leq^\square -monotonic if $\mathbf{a}_1 \leq^\square \mathbf{a}_2, \dots, \mathbf{b}_1 \leq^\square \mathbf{b}_2$ imply that $F(\mathbf{a}_1, \dots, \mathbf{b}_1) \leq^\square F(\mathbf{a}_2, \dots, \mathbf{b}_2)$.
- We say that an n -ary interval operation $F : \mathbb{I}(L) \times \dots \times \mathbb{I}(L) \rightarrow \mathbb{I}(L)$ is \leq^\diamond -monotonic if $\mathbf{a}_1 \leq^\diamond \mathbf{a}_2, \dots, \mathbf{b}_1 \leq^\diamond \mathbf{b}_2$ imply that $F(\mathbf{a}_1, \dots, \mathbf{b}_1) \leq^\diamond F(\mathbf{a}_2, \dots, \mathbf{b}_2)$.

3.3 Solution Simplified

At first glance, the above solution may seem somewhat complicated. Indeed, if we try to use the above definitions to check, e.g., whether \mathbf{a}_1 is necessarily \leq than \mathbf{a}_2 , then we will have to check *infinitely many* inequalities $a_1 \leq a_2$ for all possible pairs $a_1 \in \mathbf{a}_1$ and $a_2 \in \mathbf{a}_2$. Luckily, the above definition can be easily simplified; indeed, the following result can be easily proven:

Proposition 1.

- $\mathbf{a}_1 \leq^\square \mathbf{a}_2 \leftrightarrow a_1^+ \leq a_2^-$;
- $\mathbf{a}_1 \leq^\diamond \mathbf{a}_2 \leftrightarrow a_1^- \leq a_2^+$.

3.4 Simple to Check But Not Easy to Analyze

The above reformulation shows that both relations \leq^\square and \leq^\diamond are easy to check. However, this same result shows that these relations are not easy to analyze, because they are *not* orders [27].

Indeed, an order \leq is *reflexive* (i.e., $a \leq a$ for every a), but the relation \leq^\square is *not* reflexive: if $a^- < a^+$, then $[a^-, a^+] \not\leq^\square [a^-, a^+]$. One might suspect that \leq^\square is a *strict order*, i.e., a anti-reflexive relation (for which $a \not\leq a$ for all a), but this is not true either: for degenerate intervals, the relation is reflexive: $\mathbf{a} \leq^\square \mathbf{a}$. Similarly, the order \leq should be *transitive* (if $a \leq b$ and $b \leq c$, then $a \leq c$), but the relation \leq^\diamond is *not* transitive: e.g., $[0.9, 1.0] \leq^\diamond [0, 1]$, $[0, 1] \leq^\diamond [0, 0.1]$, but $[0.9, 1] \not\leq^\diamond [0, 0.1]$.

Since these relations are *not* orders, we cannot use standard results about monotonicity, and we therefore have to prove everything “from scratch”. This is what we will do in the next section. An interesting auxiliary question – originally formulated in [27] – is to give a complete algebraic characterization of these relations. This characterization is given in Section 4.

3.5 Additional Monotonicity Property: Inclusion Monotonicity

Let us show that, in addition to \leq^\square - and \leq^\diamond -monotonicity, it is natural to require one more monotonicity property for interval operations. Indeed, suppose that initially, we had \mathbf{a}_2 and \mathbf{b}_2 as sets of possible values of degrees of confidence in A and B . Then, by applying the interval “and”-operation $f_{\&}$, we can conclude that the degree of confidence in $A \& B$ is in $f_{\&}(\mathbf{a}_2, \mathbf{b}_2)$.

Suppose now that we have narrowed down our degrees of confidence to $\mathbf{a}_1 \subseteq \mathbf{a}_2$ and $\mathbf{b}_1 \subseteq \mathbf{b}_2$. If we apply the same interval “and”-operation to the new degrees of confidence, we get a new interval $f_{\&}(\mathbf{a}_1, \mathbf{b}_1)$. Since we have narrowed down our intervals of possible degrees of confidence, it can happen that some previously possible degrees of confidence in $A \& B$ are not possible anymore. But it is reasonable to require that if a value is now possible, then it was possible earlier as well (when we had even less knowledge about degrees of confidence). In other words, we require that every number from $f_{\&}(\mathbf{a}_1, \mathbf{b}_1)$ should belong to $f_{\&}(\mathbf{a}_2, \mathbf{b}_2)$.

In other words, we require that if $\mathbf{a}_1 \subseteq \mathbf{a}_2$ and $\mathbf{b}_1 \subseteq \mathbf{b}_2$, then $f_{\&}(\mathbf{a}_1, \mathbf{b}_1) \subseteq f_{\&}(\mathbf{a}_2, \mathbf{b}_2)$. In mathematical terms, we require that the interval “and”-operation $f_{\&}(\mathbf{a}, \mathbf{b})$ be monotonic relative to set inclusion \subseteq , i.e., in short, *inclusion monotonic*.

Definition 5. *Let n be an arbitrary positive integer. We say that an n -ary interval operation $F : \mathbb{I}(L) \times \dots \times \mathbb{I}(L) \rightarrow \mathbb{I}(L)$ is inclusion-monotonic if $\mathbf{a}_1 \subseteq \mathbf{a}_2, \dots, \mathbf{b}_1 \subseteq \mathbf{b}_2$ imply that $F(\mathbf{a}_1, \dots, \mathbf{b}_1) \subseteq F(\mathbf{a}_2, \dots, \mathbf{b}_2)$.*

Now, we are ready for the main result.

4 Main Result

Although our main interest is in binary operations over subintervals of the interval $[0, 1]$, we will formulate this result in the most general terms: as a result about operations of arbitrary arity over subintervals of an arbitrary ordered set L .

Definition 6. *Let n be an arbitrary positive integer. We say that an n -ary interval operation $F : \mathbb{I}(L) \times \dots \times \mathbb{I}(L) \rightarrow \mathbb{I}(L)$ is obtained by interval computations if there exists an n -ary \leq -monotonic function $f : L \times \dots \times L \rightarrow L$ for which*

$$F([a^-, a^+], \dots, [b^-, b^+]) = [f(a^-, \dots, b^-), f(a^+, \dots, b^+)].$$

Theorem 1.

- *Every operation F obtained by interval computations is \leq^\square -monotonic, \leq^\diamond -monotonic, and inclusion-monotonic.*
- *Every \leq^\square -monotonic, \leq^\diamond -monotonic, and inclusion-monotonic interval operation F is obtained by interval computations.*

The second part of this theorem says that *every interval-based operation which satisfies the above natural monotonicity requirement is obtained by interval computations*. Thus, for binary operations over $\mathbb{I}([0, 1])$, we did provide a fundamental justification for the traditional pragmatic approach to interval-valued operations.

Editorial Comment. For the convenience of the readers who are interested in the results but not in the technical details of the proofs, all the proofs are placed in the special Proofs section located at the end of the paper.

Technical Comment. In the second part of Theorem 1, we required that the interval operation F be both \leq^\square -monotonic and \leq^\diamond -monotonic. As one can see from the proof, it is sufficient to require that F is \leq^\square -monotonic; then \leq^\diamond -monotonicity follows automatically.

Historical Comment. A similar result was proven, in [7], under a different monotonicity assumption: that the operation F is monotonic relative to the component-wise order:

$$[a_1^-, a_1^+] \leq [a_2^-, a_2^+] \leftrightarrow (a_1^- \leq a_2^- \ \& \ a_1^+ \leq a_2^+).$$

In contrast to the relations \leq^\square and \leq^\diamond , the above relation is an order.

5 Auxiliary Results

Normally, we require that the relation \leq between degrees of certainty be an order, i.e., a relation which satisfies the following three properties:

- it is *reflexive* ($a \leq a$);
- it is *transitive* ($a \leq b$ and $b \leq c$ imply $a \leq c$); and
- it is *antisymmetric* ($a \leq b$ and $b \leq a$ imply $a = b$).

In the previous section, we mentioned that neither \leq^\square nor \leq^\diamond are orders. What are they?

In this section, we give exact algebraic characterizations of these two relations. To describe these results, let us recall the definition of a restriction of a relation to a subset. Let S be an arbitrary set, let R be an arbitrary relation on this set, and let $S' \subseteq S$ be a subset of S . Then, we define a *restriction* $R|_{S'}$ of R to S' as follows: if $a, b \in S'$ then $a R|_{S'} b$ if and only if $a R b$.

Theorem 2.

- *Let L be an arbitrary partially ordered set, and let S be an arbitrary subset of $\mathbb{I}(L)$. Then, the restriction of \leq^\square on S is transitive and antisymmetric.*
- *Let S be an arbitrary set with a transitive antisymmetric relation R . Then, there exists a partially ordered set L and a subset S' of the interval set $\mathbb{I}(L)$ such that the relation R on S is isomorphic to the restriction of \leq^\square to S' .*

Theorem 3.

- *Let L be an arbitrary partially ordered set, and let S be an arbitrary subset of $\mathbb{I}(L)$. Then, the restriction of \leq^\diamond on S is reflexive.*
- *Let S be an arbitrary set with a reflexive relation R . Then, there exists a partially ordered set L and a subset S' of the interval set $\mathbb{I}(L)$ such that the relation R on S is isomorphic to the restriction of \leq^\diamond to S' .*

So, both relations appear naturally if we divide the three properties describing order into two groups: reflexivity in one group, and transitivity and antisymmetry in another group.

- If we only keep properties from the first group, we get \leq^\diamond .
- If we only keep properties from the second group, we get \leq^\square .
- If we keep properties from both groups, we get a normal order relation.

6 Proofs

6.1 Proof of Theorem 1

1°. The first part is reasonable straightforward: if the interval operation F is obtained by interval computations from some monotonic operation

$$f : L \times \dots \times L \rightarrow L,$$

then from \leq -monotonicity of f , one can easily prove that F is \leq^\square -, \leq^\diamond -, and inclusion-monotonic.

2°. To complete the proof of the theorem, we must therefore prove its second part: that every \leq^\square -, \leq^\diamond -, and inclusion-monotonic interval operation F is obtained by interval computations.

We will actually prove this result without requiring that F is \leq^\diamond -monotonic. Then, from the first part, it will follow that \leq^\diamond -monotonicity is automatically satisfied.

So, let F be \leq^\square - and inclusion-monotonic. The result of applying F is an interval. Let us denote its lower endpoint by F^- and its upper endpoint by F^+ .

2.1°. Let us first prove that when all inputs to F are degenerate intervals, then the output is also degenerate, i.e., for every $a, \dots, b \in L$, we have

$$F^-([a, a], \dots, [b, b]) = F^+([a, a], \dots, [b, b]).$$

Indeed, by definition of \leq^\square , for every $a \in L$, we have

$$[a, a] \leq^\square [a, a].$$

So, $[a, a] \leq^\square [a, a], \dots, [b, b] \leq^\square [b, b]$, and due to \leq^\square -monotonicity of the operation F , we conclude that

$$F([a, a], \dots, [b, b]) \leq^\square F([a, a], \dots, [b, b]).$$

By definition of \leq^\square , from

$$F^-([a, a], \dots, [b, b]) \in F([a, a], \dots, [b, b])$$

and

$$F^+([a, a], \dots, [b, b]) \in F([a, a], \dots, [b, b]),$$

we can conclude that

$$F^+([a, a], \dots, [b, b]) \leq F^-([a, a], \dots, [b, b]).$$

On the other hand, since F^- and F^+ are endpoints of the interval, we have

$$F^-([a, a], \dots, [b, b]) \leq F^+([a, a], \dots, [b, b]).$$

Thus,

$$F^-([a, a], \dots, [b, b]) = F^+([a, a], \dots, [b, b]).$$

The statement is proven.

2.2°. Let us define a function $f : L \times \dots \times L \rightarrow L$ as follows: for every $a, \dots, b \in L$, we define

$$f(a, \dots, b) \stackrel{\text{def}}{=} F^-([a, a], \dots, [b, b]) = F^+([a, a], \dots, [b, b]).$$

Then, for degenerate intervals, we have

$$F([a, a], \dots, [b, b]) = [f(a, \dots, b), f(a, \dots, b)].$$

We will complete the proof of the theorem by showing two things:

- that thus defined function f is monotonic, and
- that

$$F([a^-, a^+], \dots, [b^-, b^+]) = [f(a^-, \dots, b^-), f(a^+, \dots, b^+)]$$

for all possible intervals $[a^-, a^+], \dots, [b^-, b^+] \in \mathbb{I}(L)$.

2.3°. Let us prove that the function f (defined in Part 2.2 of this proof) is monotonic. In other words, let us prove that if $a_1 \leq a_2, \dots, b_1 \leq b_2$, then $f(a_1, \dots, b_1) \leq f(a_2, \dots, b_2)$.

Indeed, let $a_1 \leq a_2, \dots, b_1 \leq b_2$. By definition of \leq^\square , we can therefore conclude that $[a_1, a_1] \leq^\square [a_2, a_2], \dots, [b_1, b_1] \leq^\square [b_2, b_2]$. Due to \leq^\square -monotonicity of the operation F , we conclude that

$$F([a_1, a_1], \dots, [b_1, b_1]) \leq^\square F([a_2, a_2], \dots, [b_2, b_2]).$$

We already know, from Part 2.3 of this proof, that

$$F([a_1, a_1], \dots, [b_1, b_1]) = [f(a_1, \dots, b_1), f(a_1, \dots, b_1)]$$

and

$$F([a_2, a_2], \dots, [b_2, b_2]) = [f(a_2, \dots, b_2), f(a_2, \dots, b_2)].$$

Thus, the above “necessarily \leq ” relation means that

$$f(a_1, \dots, b_1) \leq f(a_2, \dots, b_2).$$

The statement is proven.

2.4°. Let us now prove that

$$F([a^-, a^+], \dots, [b^-, b^+]) = [f(a^-, \dots, b^-), f(a^+, \dots, b^+)]$$

for all possible intervals $[a^-, a^+], \dots, [b^-, b^+] \in \mathbb{I}(L)$, i.e., that for all possible intervals,

$$F^-([a^-, a^+], \dots, [b^-, b^+]) = f(a^-, \dots, b^-)$$

and

$$F^+([a^-, a^+], \dots, [b^-, b^+]) = f(a^+, \dots, b^+).$$

2.4.1°. Let us first prove that

$$f(a^-, \dots, b^-) \leq F^-([a^-, a^+], \dots, [b^-, b^+]).$$

Indeed, from the definition of \leq^\square , we can easily conclude that for every interval $[a^-, a^+]$, we have $[a^-, a^-] \leq^\square [a^-, a^+]$.

From the fact that $[a^-, a^-] \leq^\square [a^-, a^+], \dots, [b^-, b^-] \leq^\square [b^-, b^+]$, and that F is \leq^\square -monotonic, we conclude that

$$F([a^-, a^-], \dots, [b^-, b^-]) \leq^\square F([a^-, a^+], \dots, [b^-, b^+]).$$

According to Proposition 1, this means that

$$F^+([a^-, a^-], \dots, [b^-, b^-]) \leq F^-([a^-, a^+], \dots, [b^-, b^+]).$$

We already know, from Part 2.2 of this proof, that

$$F^+([a^-, a^-], \dots, [b^-, b^-]) = f(a^-, \dots, b^-).$$

Thus, the above inequality is exactly what we want to prove. The statement is proven.

2.4.2°. Let us now prove that

$$F^-([a^-, a^+], \dots, [b^-, b^+]) \leq f(a^-, \dots, b^-).$$

Indeed, for each of the input intervals, we have $[a^-, a^-] = \{a^-\} \subseteq [a^-, a^+], \dots, [b^-, b^-] = \{b^-\} \subseteq [b^-, b^+]$. Since the operation F is inclusion-monotonic, we conclude that

$$\begin{aligned} F([a^-, a^-], \dots, [b^-, b^-]) &\subseteq F([a^-, a^+], \dots, [b^-, b^+]) = \\ &[F^-([a^-, a^+], \dots, [b^-, b^+]), F^+([a^-, a^+], \dots, [b^-, b^+])]. \end{aligned}$$

Due to Parts 2.1 and 2.2. of this proof, we have

$$F([a^-, a^-], \dots, [b^-, b^-]) = \{f(a^-, \dots, b^-)\}.$$

Thus, the above inclusion means that

$$f(a^-, \dots, b^-) \in [F^-([a^-, a^+], \dots, [b^-, b^+]), F^+([a^-, a^+], \dots, [b^-, b^+])].$$

By definition of an interval, this means, in particular, that

$$F^-([a^-, a^+], \dots, [b^-, b^+]) \leq f(a^-, \dots, b^-).$$

The statement is proven.

2.4.3°. From Parts 2.4.1 and 2.4.2 of this proof, we can now conclude that

$$F^-([a^-, a^+], \dots, [b^-, b^+]) = f(a^-, \dots, b^-).$$

2.4.4°. Let us now start proving the second inequality from Part 2.4 by first proving that

$$F^+([a^-, a^+], \dots, [b^-, b^+]) \leq f(a^+, \dots, b^+).$$

Indeed, from the definition of \leq^\square , we can easily conclude that for every interval $[a^-, a^+]$, we have $[a^-, a^+] \leq^\square [a^+, a^+]$.

From the fact that $[a^-, a^+] \leq^\square [a^+, a^+]$, \dots , $[b^-, b^+] \leq^\square [b^+, b^+]$, and that F is \leq^\square -monotonic, we conclude that

$$F([a^-, a^+], \dots, [b^-, b^+]) \leq^\square F([a^+, a^+], \dots, [b^+, b^+]).$$

According to Proposition 1, this means that

$$F^+([a^-, a^+], \dots, [b^-, b^+]) \leq F^-([a^+, a^+], \dots, [b^+, b^+]).$$

We already know, from Part 2.2 of this proof, that

$$F^-([a^+, a^+], \dots, [b^+, b^+]) = f(a^+, \dots, b^+).$$

Thus, the above inequality is exactly what we want to prove. The statement is proven.

2.4.5°. Let us now prove that

$$f(a^+, \dots, b^+) \leq F^+([a^-, a^+], \dots, [b^-, b^+]).$$

Indeed, for each of the input intervals, we have $[a^+, a^+] = \{a^+\} \subseteq [a^-, a^+], \dots, [b^+, b^+] = \{b^+\} \subseteq [b^-, b^+]$. Since the operation F is inclusion-monotonic, we conclude that

$$F([a^+, a^+], \dots, [b^+, b^+]) \subseteq F([a^-, a^+], \dots, [b^-, b^+]) = [F^-([a^-, a^+], \dots, [b^-, b^+]), F^+([a^-, a^+], \dots, [b^-, b^+])].$$

Due to Parts 2.1 and 2.2. of this proof, we have

$$F([a^+, a^+], \dots, [b^+, b^+]) = \{f(a^+, \dots, b^+)\}.$$

Thus, the above inclusion means that

$$f(a^+, \dots, b^+) \in [F^-([a^-, a^+], \dots, [b^-, b^+]), F^+([a^-, a^+], \dots, [b^-, b^+])].$$

By definition of an interval, this means, in particular, that

$$f(a^+, \dots, b^+) \leq F^+([a^-, a^+], \dots, [b^-, b^+]).$$

The statement is proven.

2.4.6°. From Parts 2.4.4 and 2.4.5 of this proof, we can now conclude that

$$F^+([a^-, a^+], \dots, [b^-, b^+]) = f(a^+, \dots, b^+).$$

The theorem is proven.

6.2 Proof of Theorem 2

The first part of the theorem easily follows from Proposition 1, so it is sufficient to prove the second part.

Let S be a set with a transitive antisymmetric relation R . Let

$$S_r \stackrel{\text{def}}{=} \{a \in S \mid a R a\}$$

denote the set of all reflexive elements of S , and let

$$S_i \stackrel{\text{def}}{=} \{a \in S \mid \neg a R a\}$$

denote the set of all irreflexive elements of S . Let us define L as

$$L \stackrel{\text{def}}{=} S_r \cup (S_i \times \{-, +\}),$$

i.e., as a set consisting of:

- all reflexive element of S , and

- of pairs $\langle a, - \rangle$ and $\langle a, + \rangle$, where $a \in S_i$,

and let us define the relation \leq on L as follows:

- for every $a, b \in S_r$, we have $a \leq b$ if and only if $a R b$;
- for $a \in S_r$ and $b \in S_i$, we have:
 - $a \leq \langle b, - \rangle$ if and only if $a R b$, and
 - $a \leq \langle b, + \rangle$ if and only if $a R b$;
- for $a \in S_i$ and $b \in S_r$, we have:
 - $\langle a, - \rangle \leq b$ if and only if $a R b$, and
 - $\langle a, + \rangle \leq b$ if and only if $a R b$.
- for every $a \in S_i$, $\langle a, - \rangle \leq \langle a, - \rangle$, $\langle a, - \rangle \leq \langle a, + \rangle$, and $\langle a, + \rangle \leq \langle a, + \rangle$;
- finally, for $a, b \in S_i$, $a \neq b$, we have:
 - $\langle a, - \rangle \leq \langle b, - \rangle$ if and only if $a R b$;
 - $\langle a, - \rangle \leq \langle b, + \rangle$ if and only if $a R b$;
 - $\langle a, + \rangle \leq \langle b, - \rangle$ if and only if $a R b$;
 - $\langle a, + \rangle \leq \langle b, + \rangle$ if and only if $a R b$.

One can easily check that this relation is an order.

Let us now assign, to every element $a \in S$, an interval from $\mathbb{I}(L)$. Specifically, we assign:

- to every element $a \in S_r$, a degenerate interval $[a, a] \in \mathbb{I}(L)$, and
- to every element $a \in S_i$, an interval $[\langle a, - \rangle, \langle a, + \rangle] \in \mathbb{I}(L)$.

Due to Proposition 1 and the definition of the order on L , we have the following equivalences:

- when $a, b \in S_r$, then $[a, a] \leq^\square [b, b]$ if and only if $a R b$;
- when $a \in S_r$ and $b \in S_i$, then $[a, a] \leq^\square [\langle b, - \rangle, \langle b, + \rangle]$ if and only if $a R b$;
- when $a \in S_i$ and $b \in S_r$, then $[\langle a, - \rangle, \langle a, + \rangle] \leq^\square [b, b]$ if and only if $a R b$;
- finally, when $a, b \in S_i$, then $[\langle a, - \rangle, \langle a, + \rangle] \leq^\square [\langle b, - \rangle, \langle b, + \rangle]$ if and only if $a R b$.

Thus, the original relation R on S is isomorphic to the restriction of \leq^\square to the set S' of all intervals assigned to elements of S . The theorem is proven.

6.3 Proof of Theorem 3

The first part of this theorem easily follows from the definition of \leq^\diamond , so it is sufficient to prove the second part.

Let S be a set with a reflexive relation R . Let us define L as $S \times \{-, +\}$, i.e., as the set of all pairs $\langle a, - \rangle$ and $\langle a, + \rangle$, where $a \in S$, and let us define the relation \leq on L as follows:

- for every $a \in S$, $\langle a, - \rangle \leq \langle a, - \rangle$, $\langle a, - \rangle \leq \langle a, + \rangle$, and $\langle a, + \rangle \leq \langle a, + \rangle$;
- for $a \neq b$, we have $\langle a, - \rangle \leq \langle b, + \rangle$ if and only if $a R b$;
- for every a and b , $\langle a, + \rangle \not\leq \langle b, - \rangle$.

One can easily check that this relation is an order.

Let us now assign, to every element $a \in S$, an interval $[\langle a, - \rangle, \langle a, + \rangle] \in \mathbb{I}(L)$. Due to Proposition 1 and the definition of the order on L , we have $[\langle a, - \rangle, \langle a, + \rangle] \leq^\diamond [\langle b, - \rangle, \langle b, + \rangle]$ if and only if $a R b$. Thus, the original relation R on S is isomorphic to the restriction of \leq^\diamond to the set S' of all intervals $[\langle a, - \rangle, \langle a, + \rangle]$. The theorem is proven.

Acknowledgments

This work was supported in part by NASA under cooperative agreement NCC5-209 and grant NCC 2-1232, by the Future Aerospace Science and Technology Program (FAST) Center for Structural Integrity of Aerospace Systems, effort sponsored by the Air Force Office of Scientific Research, Air Force Materiel Command, USAF, under grant number F49620-00-1-0365, by Grant No. W-00016 from the U.S.-Czech Science and Technology Joint Fund, and Grant NSF 9710940 Mexico/Conacyt.

The authors are thankful to Qiang Zuo, to Carol and Elbert Walkers, and to Piotr Wojciechowski for their encouragement and helpful discussions, and to the anonymous referees for useful comments and ideas.

References

- [1] W. Bandler and L. J. Kohout, “Unified theory of multi-valued logical operations in the light of the checklist paradigm”, *Proc. of IEEE Conference on Systems, Man, and Cybernetics*, Halifax, Nova Scotia, Oct. 1984.
- [2] D. Berleant and H. Cheng, “A Software Tool for Automatically Verified Operations on Intervals and Probability Distributions”, *Reliable Computing*, 1998, Vol. 4, No. 1, pp. 71–82.
- [3] D. Berleant and C. Goodman-Strauss, “Bounding the Results of Arithmetic Operations on Random Variables of Unknown Dependency using Intervals”, *Reliable Computing*, 1998, Vol. 4, No. 2, pp. 147–165.

- [4] S. Ferson, “Probability bounds analysis software,” *Computing in Environmental Resource Management: Proceedings of a Special Conference*, Research Triangle Park, NC, December 2-4, 1996, Air and Waste Management Association, Pittsburgh, Pennsylvania, 1996, pp. 669-678.
- [5] S. Ferson and L. Ginzburg, “Hybrid arithmetic”, In: B. M. Ayyub (ed.), *Proceedings of the ISUMA-NAFIPS’95*, IEEE Computer Society Press, Los Alamitos, California, 1995, pp. 619–623.
- [6] S. Ferson, L. Ginzburg, V. Kreinovich, and H. Schulte, “Interval Computations as a Particular Case of a General Scheme Involving Classes of Probability Distributions”, In: J. Wolff von Gudenberg and Walter Krämer (eds.), *Proceedings of the SCAN’2000/INTERVAL’2000*, Kluwer Academic/Plenum Publishers, Dordrecht, 2001 (to appear).
- [7] M. Gehrke, C. Walker, and E. Walker, “Some comments on interval-valued fuzzy sets”, *Internat. J. Intelligent Systems*, 1996, Vol. 11, pp. 751–759.
- [8] R. Giles, “Lukasiewicz logic and fuzzy set theory”, *Internat. J. Man-Machine Stud.*, 1976, Vol. 8, pp. 313–327.
- [9] R. Hammer, M. Hocks, U. Kulisch, and D. Ratz, *Numerical toolbox for verified computing. I. Basic numerical problems*, Springer-Verlag, Heidelberg, 1993.
- [10] R. B. Kearfott, *Rigorous global search: continuous problems*, Kluwer, Dordrecht, 1996.
- [11] R. B. Kearfott and V. Kreinovich (eds.), *Applications of Interval Computations*, Kluwer, Dordrecht, 1996.
- [12] G. Klir and B. Yuan, *Fuzzy Sets and Fuzzy Logic: Theory and Applications*, Prentice Hall, Upper Saddle River, New Jersey, 1995.
- [13] V. Kreinovich, G. C. Mouzouris, and H. T. Nguyen, “Fuzzy rule based modeling as a universal control tool”, In: H. T. Nguyen and M. Sugeno (eds.), *Fuzzy Systems: Modeling and Control*, Kluwer, Boston, Massachusetts, 1998, pp. 135–195.
- [14] C. H. Ling, “Representation of associative functions”, *Publ. Math. Debrecen*, 1965, Vol. 12, pp. 189–212.
- [15] K. Marriott and P. J. Stuckey, *Programming with Constraints: An Introduction*, MIT Press, Cambridge, Massachusetts, 1998.
- [16] P. S. Mostert and A. L. Shields, “On the structure of semigroups on a compact manifold with boundary”, *Ann. of Math.*, 1957, Vol. 65, pp. 117–143.

- [17] H. T. Nguyen and V. Kreinovich, “Methodology of fuzzy control: an introduction”, In: H. T. Nguyen and M. Sugeno (eds.), *Fuzzy Systems: Modeling and Control*, Kluwer, Boston, Massachusetts, 1998, pp. 19–62.
- [18] H. T. Nguyen, V. Kreinovich, and Q. Zuo, “Interval-valued degrees of belief: applications of interval computations to expert systems and intelligent control”, *International Journal of Uncertainty, Fuzziness, and Knowledge-Based Systems (IJUFKS)*, 1997, Vol. 5, No. 3, pp. 317–358.
- [19] H. T. Nguyen and E. A. Walker, *First Course in Fuzzy Logic*, CRC Press, Boca Raton, Florida, 1999.
- [20] H. Ratschek and J. Rokne, *New computer methods for global optimization*, Ellis Horwood, Chichester, 1988.
- [21] B. Schweizer and A. Sklar, *Probabilistic metric spaces*, North Holland, New York, 1983.
- [22] M. H. Smith and V. Kreinovich, “Optimal strategy of switching reasoning methods in fuzzy control”, In: H. T. Nguyen, M. Sugeno, R. Tong, and R. Yager (eds.), *Theoretical aspects of fuzzy control*, J. Wiley, New York, 1995, pp. 117–146.
- [23] E. Tsang, *Foundations of Constraint Satisfaction*, Academic Press, N.Y., 1993.
- [24] I. B. Türkşen, “Interval valued fuzzy sets based on normal forms”, *Fuzzy Sets and Systems*, 1986, Vol. 20, pp. 191–210.
- [25] P. Van Hentenryck, *Constraint Satisfaction in Logic Programming*, MIT Press, Cambridge, Massachusetts, 1989.
- [26] P. Van Hentenryck, L. Michel, and Y. Deville, *Numerica: A Modeling Language for Global Optimization*, MIT Press, Cambridge, Massachusetts, 1997.
- [27] C. Walker and E. A. Walker, private communication, 1995.
- [28] L. A. Zadeh, “Fuzzy Sets”, *Information and Control*, 1965, Vol. 8, pp. 338–353.
- [29] Q. Zuo, “Description of strictly monotonic interval AND/OR operations”, *Reliable Computing*, 1995, Supplement (Extended Abstracts of APIC’95: International Workshop on Applications of Interval Computations, El Paso, TX, Febr. 23–25, 1995), pp. 232–235.